

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
11 March 2004 (11.03.2004)

PCT

(10) International Publication Number
WO 2004/019874 A3

- (51) International Patent Classification⁷: A61K 31/34, 31/335, 31/352
- (21) International Application Number: PCT/US2003/026886
- (22) International Filing Date: 27 August 2003 (27.08.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/407,210 30 August 2002 (30.08.2002) US
- (71) Applicant (for all designated States except US): ALCON, INC. [CH/CH]; Bosch 69, P.O. Box 62, CH-6331 Hünenberg (CH).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): MAY, Jesse, A. [US/US]; 4132 Hildring Drive East, Fort Worth, TX 76109 (US). DANTANARAYANA, Anura, P. [US/US]; 5824 Bayside Drive, Fort Worth, TX 76132 (US).
- (74) Agents: COPELAND, Barry, L. et al.; Alcon Research, Ltd., R & D Counsel Q-148, 6201 South Freeway, Fort Worth, TX 76134-2099 (US).
- (81) Designated States (national): AU, BR, CA, CN, JP, KR, MX, PL, US, ZA.
- (84) Designated States (regional): European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR).
- Published:
— with international search report
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- (88) Date of publication of the international search report:
1 July 2004
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: SUBSTITUTED 5-CHROMAN-5-YL-ETHYLAMINE COMPOUNDS AND THEIR USE FOR THE TREATMENT OF GLAUCOMA

(57) Abstract: Substituted 5-chroman-5-yl-ethylamine compounds are disclosed. Also disclosed are methods for the lowering and controlling of normal or elevated intraocular pressure as well as a method for the treatment of glaucoma using compositions containing one or more of the compounds of the present invention.

WO 2004/019874 A3

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US03/26886

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 31/34, 31/335, 31/352

US CL : 514/463, 462

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/463, 462

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
Please See Continuation Sheet

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X — Y	WO 00/044737 A1 (ELI LILLY AND COMPANY, USA) 03 August 2000 (03.08.2000), see entire documents.	1-7, 16 <hr/> 8-15, 17
X — Y	Database HCAPLUS on STN Online, No. 2000:55463, SCHULZE-ALEXANDRU et al. 'Quasi-Atomistic Receptor Surrogates for the 5-HT _{2A} receptor: a 3D-QSAR Study on Hallucinogenic Substances,' abstract, Quantitative Structure-Activity Relationships, 1999, 18(6), 548-560.	1-7, 16 <hr/> 8-15, 17



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T"

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X"

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y"

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&"

document member of the same patent family

Date of the actual completion of the international search

18 April 2004 (18.04.2004)

Date of mailing of the international search report

21 MAY 2004

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US

Commissioner for Patents

P.O. Box 1450

Alexandria, Virginia 22313-1450

Facsimile No. (703) 305-3230

Authorized officer

Brian Kwon

Telephone No. 703-308-1235

INTERNATIONAL SEARCH REPORT

PCT/US03/26886

Continuation of B. FIELDS SEARCHED Item 3:

STN Online

Search terms: aminoalkylbenzofurans, 5-HT agonist, serotoninc receptor antagonist, glaucoma

BLANK PAGE

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
11 March 2004 (11.03.2004)

PCT

(10) International Publication Number
WO 2004/019874 A2

- (51) International Patent Classification⁷: A61K (US). DANTANARAYANA, Anura, P. [US/US]; 5824 Bayside Drive, Fort Worth, TX 76132 (US).
- (21) International Application Number: PCT/US2003/026886 (74) Agents: COPELAND, Barry, L. et al.; Alcon Research, Ltd., R & D Counsel Q-148, 6201 South Freeway, Fort Worth, TX 76134-2099 (US).
- (22) International Filing Date: 27 August 2003 (27.08.2003)
- (25) Filing Language: English (81) Designated States (*national*): AU, BR, CA, CN, JP, KR, MX, PL, US, ZA.
- (26) Publication Language: English (84) Designated States (*regional*): European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR).
- (30) Priority Data: 60/407,210 30 August 2002 (30.08.2002) US
- (71) Applicant (*for all designated States except US*): ALCON, INC. [CH/CH]; Bosch 69, P.O. Box 62, CH-6331 Hünenberg (CH). Published:
— without international search report and to be republished upon receipt of that report
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): MAY, Jesse, A. [US/US]; 4132 Hildring Drive East, Fort Worth, TX 76109

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: SUBSTITUTED 5-CHROMAN-5-YL-ETHYLAMINE COMPOUNDS AND THEIR USE FOR THE TREATMENT OF GLAUCOMA

(57) Abstract: Substituted 5-chroman-5-yl-ethylamine compounds are disclosed. Also disclosed are methods for the lowering and controlling of normal or elevated intraocular pressure as well as a method for the treatment of glaucoma using compositions containing one or more of the compounds of the present invention.

WO 2004/019874 A2

- 1 -

**SUBSTITUTED 5-CHROMAN-5-YL-ETHYLAMINE COMPOUNDS AND THEIR
USE FOR THE TREATMENT OF GLAUCOMA**

5 BACKGROUND OF THE INVENTION

The present invention relates to substituted 5-chroman-5-yl-ethylamine compounds. These novel compounds are useful for lowering and controlling normal or elevated intraocular pressure (IOP) and for treating glaucoma.

10 The disease state referred to as glaucoma is characterized by a permanent loss of visual function due to irreversible damage to the optic nerve. The several morphologically or functionally distinct types of glaucoma are typically characterized by elevated IOP, which is considered to be causally related to the pathological course of the disease. Ocular hypertension is a condition wherein intraocular pressure is elevated but no apparent loss of
15 visual function has occurred; such patients are considered to be a high risk for the eventual development of the visual loss associated with glaucoma. If glaucoma or ocular hypertension is detected early and treated promptly with medications that effectively reduce elevated intraocular pressure, loss of visual function or its progressive deterioration can generally be ameliorated. Drug therapies that have proven to be effective for the reduction of intraocular
20 pressure include both agents that decrease aqueous humor production and agents that increase the outflow facility. Such therapies are in general administered by one of two possible routes, topically (direct application to the eye) or orally.

There are some individuals who do not respond well when treated with certain existing glaucoma therapies. There is, therefore, a need for other topical therapeutic agents
25 that control IOP.

Serotonergic 5-HT_{1A} agonists have been reported as being neuroprotective in animal models and many of these agents have been evaluated for the treatment of acute stroke among

other indications. This class of compounds has been mentioned for the treatment of glaucoma (lowering and controlling IOP), see e.g., WO 98/18458 and EP 0771563A2. Osborne *et al.* (1996) teach that 8-hydroxy-2-(di-*n*-propylamino)tetralin (8-OH-DPAT) (a 5-HT_{1A} agonist) reduces IOP in rabbits. Wang *et al.* (1997 and 1998) indicate that 5-methylurapidil, an α_{1A} antagonist and 5-HT_{1A} agonist lowers IOP in the monkey, but due to its α_{1A} receptor activity. Also, 5-HT_{1A} antagonists are disclosed as being useful for the treatment of glaucoma (elevated IOP) (e.g., WO 92/0338). Furthermore, WO 97/35579 and U.S. 5,578,612 relate to the use of 5-HT₁ and 5-HT_{1-like} agonists for the treatment of glaucoma (elevated IOP). These anti-migraine compounds are 5-HT_{1B,D,E,F} agonists, e.g., sumatriptan and naratriptan and related compounds.

It has been found that serotonergic compounds which possess agonist activity at 5-HT₂ receptors effectively lower and control normal and elevated IOP and are useful for treating glaucoma, see commonly owned co-pending application, PCT/US99/19888, incorporated in its entirety by reference herein. Compounds that act as agonists at 5-HT₂ receptors are well known and have shown a variety of utilities, primarily for disorders or conditions associated with the central nervous system (CNS). U.S. Patent No. 5,494,928 relates to certain 2-(indol-1-yl)-ethylamine derivatives that are 5-HT_{2C} agonists for the treatment of obsessive compulsive disorder and other CNS derived personality disorders. U.S. Patent No. 5,571,833 relates to tryptamine derivatives that are 5-HT₂ agonists for the treatment of portal hypertension and migraine. U.S. Patent No. 5,874,477 relates to a method for treating malaria using 5-HT_{2A/2C} agonists. U.S. Patent No. 5,902,815 relates to the use of 5-HT_{2A} agonists to prevent adverse effects of NMDA receptor hypo-function. WO 98/31354 relates to 5-HT_{2B} agonists for the treatment of depression and other CNS conditions. U.S. Patent 6,380,238 and International Patent Applications WO 01/12602 and WO 00/44753

relate to indoline derivatives and U.S. Patents 6,433,175 and 6,365,598 relate to certain indole derivatives as 5-HT_{2B} and 5-HT_{2C} receptor agonists for the treatment of a variety of disorders of the central nervous system, but especially for the treatment of obesity. WO 00/35922 relates to certain pyrazino[1,2-a]quinoxaline derivatives as 5-HT_{2C} agonists for the treatment of
5 obsessive compulsive disorder, depression, eating disorders, and other disorders involving the CNS. WO 00/77002 and WO 00/77010 relate to certain substituted tetracyclic pyrido[4,3-b]indoles as 5-HT_{2C} agonists with utility for the treatment of central nervous system disorders including obesity, anxiety, depression, sleep disorders, cephalic pain, and social phobias among others. Agonist response at the 5-HT_{2A} receptor is reported to be the
10 primary activity responsible for hallucinogenic activity, with some lesser involvement of the 5-HT_{2C} receptor possible (Fiorella *et al.* 1995).

5-Hydroxytryptamine (serotonin) does not cross the blood-brain barrier and enter the brain. However, in order to increase brain serotonin levels the administration of 5-hydroxytryptophane can be employed. The transport of 5-hydroxy-tryptophane into the brain readily
15 occurs, and once in the brain 5-hydroxy-tryptophane is rapidly decarboxylated to provide serotonin. Since the treatment of glaucoma is preferably with compounds that do not enter the CNS, relatively polar compounds that are 5-HT₂ agonists and have incorporated into their structure a phenolic hydroxyl group that can be considered comparable to that of serotonin, are of particular interest.

20 2-(6,7-Dimethoxy-2,3-dihydro-benzofuran-4-yl)-ethylamine has been synthesized and shown to have affinity for 5-HT₂ receptors (Monte *et al.* 1997).

The preparation of 2-(7-bromo-5-methoxy-2,3-dihydro-benzofuran-4-yl)-1-methylethylamine has been reported (Waldman *et al.* 1996) and this compound has been shown to have a high affinity for the 5-HT_{2A} receptor and to generalize to LSD in drug

discrimination studies (Nichols *et al.* 1991). Similarly, substituted 5-(2-aminopropyl)-benzodifurans and substituted 5-(2-aminopropyl)-benzodipyranes have been shown to have high affinity for 5-HT₂ receptors and to generalize to LSD in drug discrimination studies (Monte *et al.* 1996; Parker *et al.* 1998; Whiteside *et al.* 2002).

5 Benzofurans, such as 2-benzofuran-4-yl-1-methylethylamine and related compounds, are reported to have agonist activity at the 5-HT_{2C} receptor and thereby be useful for the treatment of a variety of central nervous system disorders, such as seizure and eating disorders among others (WO 00/44737).

Amides of substituted 3-chroman-5-yl-alkylamines and 3-(2,3-dihydro-benzofuran-
10 4-yl)-alkylamines have agonist or antagonist activity at melatonin receptors and thereby are useful in the treatment of disorders regulated by melatonin. These include chronobiological disorders such as seasonal affective disorders and insomnia, or psychiatric disorders such as bipolar disorders and depression (U. S. 5,981,572).

Accordingly, there is a need to provide new compounds which avoid the
15 disadvantages described above and which provide increased chemical stability and a desired length of therapeutic activity, for instance, in decreasing intraocular pressure and treating glaucoma.

SUMMARY OF THE PRESENT INVENTION


20 A feature of the present invention is to provide novel compounds which are 5-HT₂ agonists.

Another feature of the present invention is to provide compounds which have increased chemical stability and which are useful in lowering and controlling normal or elevated intraocular pressure and/or treating glaucoma.

Another feature of the present invention is to provide compounds which provide a desired level of therapeutic activity in lowering and controlling normal or elevated intraocular pressure and/or treating glaucoma.

Additional features and advantages of the present invention will be set forth in part in the description that follows, and in part will be apparent from the description, or may be learned by practice of the present invention. The objectives and other advantages of the present invention will be realized and attained by means of the elements and combinations particularly pointed out in the description and appended claims.

To achieve these and other advantages, and in accordance with the purposes of the present invention, as embodied and broadly described herein, the present invention relates to a compound having the Formula I:



In this formula, R^1 is hydrogen or an alkyl group, such as C_{1-4} alkyl;
 R^2 is hydrogen, an alkyl group such as C_{1-4} alkyl, or R^1 and R^2 can together be $(CH_2)_{2-4}$ to complete a heterocyclic ring;
 R^3 is hydrogen, hydroxyl, an alkoxy group such as C_{1-4} alkoxy, or halogen;
 R^4 and R^5 are independently selected from hydrogen, halogen, nitrile, an alkoxy group such as C_{1-4} alkoxy, an alkylthio such as C_{1-6} alkylthiol, an alkyl group such as C_{1-4} alkyl, a substituted alkyl group such as C_{1-4} alkyl substituted with halogen or C_{1-6} alkoxy; or R^4 and R^5 can together be $(CH_2)_m$ to complete a cycloalkyl ring, or they can together complete a phenyl or thiophene ring which can be unsubstituted or substituted with halogen, an alkyl

group such as C₁₋₄alkyl or an alkoxy group such as C₁₋₄alkoxy;

m = 3 - 4;

n = 0 - 2;

R⁶ is hydrogen, hydroxyl, an alkoxy group such as C₁₋₄alkoxy, a substituted alkoxy such as
5 C₁₋₄alkoxy substituted with hydroxyl, halogen, or NR⁷N⁸, or OC(=O)C₁₋₆alkyl, =O, NR⁷R⁸,
an alkyl group such as C₁₋₄alkyl, a substituted alkyl group such as C₁₋₄alkyl substituted with
hydroxyl, halogen, or NR⁷R⁸, however, when n = 0, R⁶ cannot be hydrogen;

X is an alkoxy group such as C₁₋₄alkoxy, hydroxyl, or halogen;

R⁷ and R⁸ are independently selected from hydrogen, an alkyl group such as C₁₋₄alkyl, or
10 C(=O)C₁₋₆alkyl; and the dashed bond denotes a single or double bond.

Pharmaceutically acceptable salts and solvates of Formula I are also part of the present
invention.

The present invention further relates to pharmaceutical compositions containing at
least one compound of Formula I.

15 The present invention further relates to methods to lower and/or control normal or
elevated intraocular pressure by administering an effective amount of a composition
containing a compound having Formula I as described above.


The present invention also relates to a method for treating glaucoma which involves
administering an effective amount of a composition containing a compound having Formula I
20 as described above.

It is to be understood that both the foregoing general description and the following
detailed description are exemplary and explanatory only and are intended to provide a further
explanation of the present invention, as claimed.

DETAILED DESCRIPTION OF THE PRESENT INVENTION

The present invention relates to a variety of compounds which are useful according to the present invention. These compounds are generally represented by the following Formula I.

Formula I



In this formula, R^1 is hydrogen or an alkyl group, such as C_{1-4} alkyl;

R^2 is hydrogen, an alkyl group such as C_{1-4} alkyl, or R^1 and R^2 can together be $(CH_2)_{2-4}$ to complete a heterocyclic ring;

R^3 is hydrogen, hydroxyl, an alkoxy group such as C_{1-4} alkoxy, or halogen;

R^4 and R^5 are independently selected from hydrogen, halogen, nitrile, an alkoxy group such as C_{1-4} alkoxy, an alkylthio such as C_{1-6} alkylthiol, an alkyl group such as C_{1-4} alkyl, a substituted alkyl group such as C_{1-4} alkyl substituted with halogen or C_{1-6} alkoxy; or R^4 and R^5 can together be $(CH_2)_m$ to complete a cycloalkyl ring, or they can together complete a phenyl or thiophene ring which can be unsubstituted or substituted with halogen, an alkyl group such as C_{1-4} alkyl or an alkoxy group such as C_{1-4} alkoxy;

$m = 3 - 4$;

$n = 0 - 2$;

R^6 is hydrogen, hydroxyl, an alkoxy group such as C_{1-4} alkoxy, a substituted alkoxy group such as C_{1-4} alkoxy substituted with hydroxyl, halogen, or NR^7R^8 , $OC(=O)C_{1-6}$ alkyl, $=O$, NR^7R^8 , an alkyl group such as C_{1-4} alkyl, or a substituted alkyl group such as C_{1-4} alkyl

substituted with hydroxyl, halogen, or NR^7R^8 , however, when $n = 0$, R^6 cannot be hydrogen;

X is an alkoxy group such as C_{1-4} alkoxy, hydroxyl, or halogen;

R^7 and R^8 are independently selected from hydrogen, an alkyl group such as C_{1-4} alkyl, or

5 $\text{C}(=\text{O})\text{C}_{1-6}$ alkyl; and the dashed bond denotes a single or double bond.

Pharmaceutically acceptable salts and solvates of Formula I are also part of the present invention.

Preferred compounds are:

Wherein R^1 and R^3 are hydrogen;

10 R^2 is C_{1-4} alkyl;

R^4 and R^5 are independently selected from halogen, nitrile, C_{1-4} alkoxy, C_{1-6} alkylthiol, C_{1-4} alkyl, C_{1-4} alkyl substituted with halogen, or R^4 and R^5 can together be $(\text{CH}_2)_m$ to complete a cycloalkyl ring, or they can together complete a phenyl or thiophene ring which can be unsubstituted or substituted with halogen, C_{1-4} alkyl;

15 $m = 3 - 4$;

$n = 1$;

R^6 is hydroxyl, C_{1-4} alkoxy, C_{1-4} alkoxy substituted with hydroxyl, halogen, or NR^7R^8 , $\text{OC}(=\text{O})\text{C}_{1-6}$ alkyl, NR^7R^8 , or C_{1-4} alkyl substituted with hydroxyl, halogen, or NR^7R^8 , however, when $n = 0$, R^6 cannot be hydrogen;

20 X is C_{1-4} alkoxy or hydroxyl;

R^7 and R^8 are independently selected from hydrogen, C_{1-4} alkyl, or $\text{C}(=\text{O})\text{C}_{1-6}$ alkyl.

Representative examples of preferred novel compounds of Formula I are:

5-(2-Aminopropyl)-8-bromo-6-methoxy-chroman-3-ol;

5-((R)-2-Aminopropyl)-8-bromo-6-methoxy-chroman-3-ol;

5-(2-Aminopropyl)-6-methoxy-8-methyl-chroman-3-ol;

5-((R)-2-Amino-1-hydroxy-propyl)-8-bromo-6-methoxy-chroman-3-ol;

Cyclopropanecarboxylic acid 5-((R)-2-aminopropyl)-8-bromo-6-methoxy-chroman-3-yl ester;

5 [5-(2-Aminopropyl)-6-methoxy-8-methyl-chroman-3-yl]-methanol;

5-(2-Aminopropyl)-8-iodo-chroman-3,6-diol; or

[4-(2-Aminopropyl)-5-methoxy-7-methyl-2,3-dihydro-benzofuran-2-yl]-methanol;

or combinations thereof.

It is recognized that compounds of Formula I can contain one or more chiral centers.


10 This invention contemplates all enantiomers, diastereomers and, mixtures thereof.

In the above definitions, the total number of carbon atoms in a substituent group is indicated by the C_{i,j} prefix where the numbers i and j define the number of carbon atoms; this definition includes straight chain, branched chain, and cyclic alkyl or (cyclic alkyl)alkyl groups. A substituent may be present either singly or multiply when incorporated into the indicated structural unit. For example, the substituent halogen, which means fluorine, chlorine, bromine, or iodine, would indicate that the unit to which it is attached may be substituted with one or more halogen atoms, which may be the same or different.

SYNTHESIS


20 The compounds of Formula I can be prepared by using one of several synthetic procedures. For example, compounds of Formula I where n is 1 can be prepared from the appropriately substituted chromanols 1 as described in Scheme 1 (Chambers *et al.* 2001)].

Scheme 1




Other compounds of Formula I can be prepared from 4 through selected functional group transformations well known in the art. For example, initial protection of the primary amine group followed by activation of the hydroxyl group by formation of a sulfonate ester, e.g. methansulfonyl, and subsequent reaction with a desired nucleophile such as alkylamines, dialkylamines, aryl or alkylthiols, and the like, will provide compounds 6 of Formula I (Scheme 2). Furthermore, direct oxidation of 4 with a suitable oxidizing agent, for example, a hypervalent iodine reagent, such as *o*-iodoxybenzoic acid (Frigerio *et al.* 1995), provides the ketone 8, which can be functionalized to provide yet other compounds of Formula I, such as 9, via reductive alkylation, and 7, via Grignard addition.

Scheme 2




The desired chromanols 4 can be prepared from the appropriately substituted ortho-bromophenols 10, which can be purchased from commercial sources or prepared by known procedures, as described in Scheme 3. Reaction of phenols 10 with epibromohydrin using any of a variety of well known alkylation conditions, such as DMF/NaH, provides the intermediate epoxide 11. It can be advantageous to directly effect cyclization of 11 to compound 4 by treatment with a suitable base, such as *n*-butyllithium or under Grignard conditions. Alternately, depending on the specific substituents present, it can be more advantageous to convert 11 initially to the protected haloether 13, which will more readily undergo the cyclization reaction to provide compounds 4.

Scheme 3




A specific example of Formula I is the preparation of the compound 21, wherein n is 1, R^1 , R^3 , and R^4 are hydrogen, R^2 is methyl, X is bromo, and R^6 is hydroxy. This compound can be prepared from 2,6-dibromo-4-methoxyphenol (Curran *et al.* 1996) as described in Scheme 4.

Scheme 4



- 5 Alternately, compounds of Formula I can be prepared from propargyloxy substituted intermediates (25) via initial Claisen rearrangement reactions (Plug *et al.* 1992; Macor *et al.* 1994; Macor *et al.* 2000) to give the intermediate chromenes 26. Further synthetic manipulation of 26, as outlined in Scheme 5, using well-known functional group transformations provides yet other desirable compounds of Formula I.

Scheme 5



The compounds of the present invention can be used to lower and control IOP including IOP associated with normotension glaucoma, ocular hypertension, and glaucoma in warm blooded animals including humans. The compounds are preferably formulated in pharmaceutical compositions which are preferably suitable for topical delivery to the eye of the patient.

The compounds of this invention, Formula I, can be incorporated into various types of ophthalmic formulations for delivery to the eye (e.g., topically, intracamerally, or via an implant). The compounds are preferably incorporated into topical ophthalmic formulations for delivery to the eye. The compounds may be combined with ophthalmologically acceptable preservatives, viscosity enhancers, penetration enhancers, buffers, sodium chloride, and water to form an aqueous, sterile ophthalmic suspension or solution. Ophthalmic solution formulations may be prepared by dissolving a compound in a

physiologically acceptable isotonic aqueous buffer. Further, the ophthalmic solution may include an ophthalmologically acceptable surfactant to assist in dissolving the compound. Furthermore, the ophthalmic solution may contain an agent to increase viscosity, such as hydroxymethylcellulose, hydroxyethylcellulose, hydroxypropylmethylcellulose, methylcellulose, polyvinylpyrrolidone, or the like, to improve the retention of the formulation in the conjunctival sac. Gelling agents can also be used, including, but not limited to, gellan and xanthan gum. In order to prepare sterile ophthalmic ointment formulations, the active ingredient is combined with a preservative in an appropriate vehicle, such as, mineral oil, liquid lanolin, or white petrolatum. Sterile ophthalmic gel formulations may be prepared by suspending the active ingredient in a hydrophilic base prepared from the combination of, for example, carbopol-974, or the like, according to the published formulations for analogous ophthalmic preparations; preservatives and tonicity agents can be incorporated.

The compounds are preferably formulated as topical ophthalmic suspensions or solutions, with a pH of about 5 to 8. The compounds will normally be contained in these formulations in an amount 0.01% to 5% by weight, but preferably in an amount of 0.25% to 2% by weight. Thus, for topical presentation 1 to 2 drops of these formulations would be delivered to the surface of the eye 1 to 4 times per day according to the discretion of a skilled clinician.

The compounds can also be used in combination with other agents for treating glaucoma, such as, but not limited to, β -blockers (e.g., timolol, betaxolol, levobetaxolol, carteolol, levobunolol, propranolol), carbonic anhydrase inhibitors (e.g., brinzolamide and dorzolamide), α_1 antagonists (e.g., nipradolol), α_2 agonists (e.g. iopidine and brimonidine), miotics (e.g., pilocarpine and epinephrine), prostaglandin analogs (e.g., latanoprost, travoprost, unoprostone, and compounds set forth in U.S. Patent Nos. 5,889,052; 5,296,504;

5,422,368; and 5,151,444, "hypotensive lipids" (e.g., bimatoprost and compounds set forth in 5,352,708), and neuroprotectants (e.g., compounds from U.S. Patent No. 4,690,931, particularly eliprodil and R-eliprodil, as set forth in WO 01/85152, and appropriate compounds from WO 94/13275, including memantine.

5 In the formulas described above, the alkyl group can be straight-chain, branched or cyclic and the like. Halogen includes Cl, Br, F, or I. Alkoxy is understood as an alkyl group bonded through an oxygen atom.

The compounds of the present invention preferably function as 5-HT₂ agonists and preferably do not enter the CNS. In more detail, the particular compounds of the present
10 invention have incorporated into their structure a phenolic hydroxyl group which is considered comparable to that of serotonin and thus the compounds of the present invention preferably does not cross the blood-brain barrier and enter the brain. Compounds having the ability to be a 5-HT₂ agonist are beneficial for controlling IOP as well as the treatment of glaucoma as shown in WO 00/16761, incorporated in its entirety by reference herein.

15 The compounds of the present invention preferably provide increased chemical stability and preferably achieve the desired level of therapeutic activity which includes a lowering or controlling of IOP.

The compounds of the present invention can be used in controlling or lowering IOP in warm-blooded animals including humans. Preferably, an effective amount of the
20 compound is administered to the patient such that the IOP is controlled or lowered to acceptable levels. Furthermore, the compounds of the present invention can be used to treat glaucoma in warm-blooded animals, including humans, by administering an effective amount of the compound to a patient in need of such treatment to treat the glaucoma.

Other embodiments of the present invention will be apparent to those skilled in the art

from consideration of the present specification and practice of the present invention disclosed herein. It is intended that the present specification and examples be considered as exemplary only with a true scope and spirit of the invention being indicated by the following claims and equivalents thereof.

5

METHOD 1

5-HT₂ Receptor Binding Assay

To determine the affinities of serotonergic compounds at the 5-HT₂ receptors, their ability to compete for the binding of the agonist radioligand [¹²⁵I]DOI to brain 5-HT₂ receptors is determined as described below with minor modification of the literature procedure (Johnson *et al.* 1987)]. Aliquots of post mortem rat or human cerebral cortex homogenates (400 µL) dispersed in 50 mM Tris-HCl buffer (pH 7.4) are incubated with [¹²⁵I]DOI (80 pM final) in the absence or presence of methiothepin (10 µM final) to define total and non-specific binding, respectively, in a total volume of 0.5 mL. The assay mixture is incubated for 1 hour at 23°C in polypropylene tubes and the assays terminated by rapid vacuum filtration over Whatman GF/B glass fiber filters previously soaked in 0.3% polyethyleneimine using ice-cold buffer. Test compounds (at different concentrations) are substituted for methiothepin. Filter-bound radioactivity is determined by scintillation spectrometry on a beta counter. The data are analyzed using a non-linear, iterative curve-fitting computer program (Bowen *et al.* 1995) to determine the compound affinity parameter. The concentration of the compound needed to inhibit the [¹²⁵I]DOI binding by 50% of the maximum is termed the IC₅₀ value.

METHOD 2

5-HT₂ functional Assay: [Ca²⁺]_i Mobilization

The receptor-mediated mobilization on intracellular calcium ([Ca²⁺]_i) was studied using the Fluorescence Imaging Plate Reader (FLIPR) instrument. Rat vascular smooth muscle cells, A7r5, were grown in a normal media of DMEM / 10% FBS and 10 µg/mL gentamycin. Confluent cell monolayers were trypsinized, pelleted, and re-suspended in normal media. Cells were seeded in a 50 µL volume at a density of 20,000 cells / well in a black wall, 96-well tissue culture plate and grown for 2 days.

On the day of the experiment, one vial of FLIPR Calcium Assay Kit dye was re-suspended in 50 mL of a FLIPR buffer consisting of Hank's Balanced Salt Solution (HBSS), 20 mM HEPES, and 2.5 mM probenecid, pH 7.4. Cells were loaded with the calcium-sensitive dye by addition of an equal volume (50 µL) to each well of the 96-well plate and incubated with dye for 1h at 23°C.

Typically, test compounds were stored at 25 µM in 50% DMSO/50% Ethanol solvent. Compounds were diluted 1:50 in 20% DMSO/20% Ethanol. For "hit" screening, compounds were further diluted 1:10 in FLIPR buffer and tested at a final concentration of 10 µM. For dose-response experiments, compounds were diluted 1:50 in FLIPR buffer and serially diluted 1:10 to give a 5- or 8- point dose-response curve.

The compound plate and cell plate were placed in the FLIPR instrument. At the beginning of an experimental run, a signal test was performed to check the basal fluorescence signal from the dye-loaded cells and the uniformity of the signal across the plate. The basal fluorescence was adjusted between 8000-12000 counts by modifying the exposure time, the camera F-stop, or the laser power. Instrument settings for a typical assay were the following: laser power 0.3-0.6 W, camera F-stop F/2, and exposure time 0.4

sec. An aliquot (25 μ L) of the test compound was added to the existing 100 μ L dye-loaded cells at a dispensing speed of 50 μ L/sec. Fluorescence data were collected in real-time at 1.0 sec intervals for the first 60 secs and at 6.0 sec intervals for an additional 120 secs. Responses were measured as peak fluorescence intensity minus basal and where appropriate were expressed as a percentage of a maximum 5-HT-induced response. When the compounds were tested as antagonists against 10 μ M 5-HT, they were incubated with the cells for 15 minutes prior to the addition of 5-HT.

The above procedures were used to generate the data shown in Table 1.

Table 1

5-HT_{2A} Receptor Binding and Functional Data

Compound	IC ₅₀ , nM	EC ₅₀ , nM	Efficacy (E _{max} , %)
26	0.31	108	23
DOI	0.33	30.2	31

All of the compositions and/or methods disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the compositions and methods of this invention have been described in terms of preferred embodiments, it will be apparent to those of skill in the art that variations may be applied to the compositions and/or methods and in the steps or in the sequence of steps of the method described herein without departing from the concept, spirit and scope of the invention. More specifically, it will be apparent that certain agents which are both chemically and structurally related may be substituted for the agents described herein to achieve similar results. All such substitutions and modifications apparent to those skilled in the art are deemed to be within the spirit, scope and concept of the invention as defined by the appended claims.

References

The following references, to the extent that they provide exemplary procedural or other details supplementary to those set forth herein, are specifically incorporated herein by reference.

United States Patents and Applications

5 4,690,931

5,151,444

5,296,504

5,352,708

5,422,368

10 5,494,928

5,571,833

5,578,612

5,874,477

5,889,052

15 5,902,815

5,981,572

6,365,598

6,380,238

6,433,175

20 **Foreign Patents and Published Patent Applications**

EP 0771563 A2

PCT/US99/19888

WO 01/85152

WO 92/0338

WO 94/13275

WO 97/35579

WO 98/18458

WO 98/31354

5 WO 00/16761

WO 00/35922

WO 00/44737

WO 00/44753

WO 00/77002

10 WO 00/77010

WO 01/12602

Other Publications

W. P. Bowen *et al.*, *Trends Pharmacol. Sci.* 16:413 (1995).

15 J. J. Chambers *et al.*, *J. Med. Chem.* 44:1003 (2001).

D. Curran *et al.*, *J. Amer. Chem. Soc.* 118:3142 (1996).

D. Fiorella *et al.*, *Psychopharmacology* 121:357 (1995).

M. Frigerio *et al.*, *J. Org. Chem.* 60:7272 (1995).

M. P. Johnson *et al.*, *Neuropharmacology* 26:1803 (1987).

20 J. E. Macor *et al.*, *Tetrahedron Lett.* 35:45 (1994).

J. E. Macor, *Tetrahedron Lett.* 41:3541 (2000).

A. P. Monte *et al.*, *J. Med. Chem.* 39:2953 (1996).

A. P. Monte *et al.*, *J. Med. Chem.* 40:2997 (1997).

D. E. Nichols *et al.*, *J. Med. Chem.* 34:276 (1991).

M. A. Parker *et al.*, *J. Med. Chem.* 41:5148 (1998).

J. P. M. Plug *et al.*, *Tetrahedron Lett.* 33:2179 (1992).

Osborne *et al.*, *Ophthalmologia* 210:308-314 (1996).

S. R. Waldman *et al.*, *Tetrahedron Lett.* 37:7889 (1996).


5 Wang *et al.*, *Current Eye Research* 16(8):769-775 (1997).

Wang *et al.*, *IVOS* 39(4):S488 (1998).

M. S. Whiteside *et al.*, *Bioorg. Med. Chem.* 10:3301 (2002).

WHAT IS CLAIMED IS:

1. A compound represented by Formula I:



wherein R^1 is hydrogen or an alkyl group;

- 5 R^2 is hydrogen, an alkyl group, or R^1 and R^2 represent $(CH_2)_{2-4}$ to complete a heterocyclic ring;

R^3 is hydrogen, hydroxyl, an alkoxy group, or halogen;

- R^4 and R^5 are independently selected from hydrogen, halogen, nitrile, an alkoxy group, an alkylthiol, a substituted or unsubstituted alkyl group, or R^4 and R^5 represent $(CH_2)_m$ to
 10 complete a cycloalkyl ring, or R^4 and R^5 represent or complete a phenyl or thiophene ring which is unsubstituted or substituted with halogen, an alkyl group, or an alkoxy group;

$m = 3 - 4$;

$n = 0 - 2$;

- R^6 is hydrogen, hydroxyl, an alkoxy group, alkoxy substituted with hydroxyl, halogen, or
 15 NR^7R^8 , $OC(=O)alkyl$, $=O$, NR^7R^8 , or a substituted or unsubstituted alkyl group, wherein when $n = 0$, R^6 is not hydrogen;

X is an alkoxy group, hydroxyl or halogen;

R^7 and R^8 are independently selected from hydrogen, an alkyl group, or $C(=O)alkyl$;

or pharmaceutically acceptable salts or solvates thereof.

20

2. The compound of claim 1, wherein R^1 is hydrogen or $C_{1-4}alkyl$;

R^2 is hydrogen, C_{1-4} alkyl, or R^1 and R^2 represent $(CH_2)_{2-4}$ to complete a heterocyclic ring;

R^3 is hydrogen, hydroxyl, C_{1-4} alkoxy, or halogen;

R^4 and R^5 are independently selected from hydrogen, halogen, nitrile, C_{1-4} alkoxy, C_{1-6} alkylthiol, C_{1-4} alkyl, C_{1-4} alkyl substituted with halogen or C_{1-6} alkoxy, or R^4 and R^5 represent $(CH_2)_m$ to complete a cycloalkyl ring, or R^4 and R^5 together complete a phenyl or thiophene ring which is unsubstituted or substituted with halogen, C_{1-4} alkyl, or C_{1-4} alkoxy;

$m = 3 - 4$;

$n = 0 - 2$;

R^6 is hydrogen, hydroxyl, C_{1-4} alkoxy, C_{1-4} alkoxy substituted with hydroxyl, halogen, or NR^7R^8 , $OC(=O)C_{1-6}$ alkyl, $=O$, NR^7R^8 , C_{1-4} alkyl, or C_{1-4} alkyl substituted with hydroxyl, halogen, or NR^7R^8 , wherein when $n = 0$, R^6 is not hydrogen;

X is C_{1-4} alkoxy or hydroxyl;

R^7 and R^8 are independently selected from hydrogen, C_{1-4} alkyl, or $C(=O)C_{1-6}$ alkyl;

or pharmaceutically acceptable salts or solvates thereof.

15 3. The compound of claim 1, wherein said R^2 is hydrogen or C_{1-4} alkyl.

4. The compound of claim 1, wherein R^1 and R^3 are hydrogen;

R^2 is C_{1-4} alkyl;

R^4 and R^5 are independently selected from halogen, nitrile, C_{1-4} alkoxy, C_{1-6} alkylthiol, C_{1-4} alkyl, C_{1-4} alkyl substituted with halogen, or R^4 and R^5 together represent $(CH_2)_m$ to complete a cycloalkyl ring, or R^4 and R^5 together complete a phenyl or thiophene ring which is unsubstituted or substituted with halogen, C_{1-4} alkyl;

$m = 3 - 4$;

$n = 1$;

R^6 is hydroxyl, C_{1-4} alkoxy, C_{1-4} alkoxy substituted with hydroxyl, halogen, or

NR^7R^8 , $\text{OC}(=\text{O})\text{C}_{1-6}\text{alkyl}$, NR^7R^8 , or $\text{C}_{1-4}\text{alkyl}$ substituted with hydroxyl, halogen, or NR^7R^8 , wherein when $n = 0$, R^6 is not hydrogen;

X is $\text{C}_{1-4}\text{alkoxy}$ or hydroxyl;

R^7 and R^8 are independently selected from hydrogen, $\text{C}_{1-4}\text{alkyl}$, or $\text{C}(=\text{O})\text{C}_{1-6}\text{alkyl}$.

5 5. The compound of claim 1, wherein said compound is:

5-(2-Aminopropyl)-8-bromo-6-methoxy-chroman-3-ol;

5-((R)-2-Aminopropyl)-8-bromo-6-methoxy-chroman-3-ol;

5-(2-Aminopropyl)-6-methoxy-8-methyl-chroman-3-ol;

5-((R)-2-Amino-1-hydroxy-propyl)-8-bromo-6-methoxy-chroman-3-ol;

10 Cyclopropanecarboxylic acid 5-((R)-2-aminopropyl)-8-bromo-6-methoxy-chroman-3-yl ester;

[5-(2-Aminopropyl)-6-methoxy-8-methyl-chroman-3-yl]-methanol;

5-(2-Aminopropyl)-8-iodo-chroman-3,6-diol; or

[4-(2-Aminopropyl)-5-methoxy-7-methyl-2,3-dihydro-benzofuran-2-yl]-methanol;

15 or combinations thereof.

6. The compound of claim 1, wherein said X is hydroxyl.

7. The compound of claim 1, wherein said X is an alkoxy group.

8. A method of controlling normal or elevated intraocular pressure comprising administering a pharmaceutically effective amount of a composition comprising at least one
20 compound of claim 1.

9. The method of claim 8, wherein R^1 is hydrogen or $\text{C}_{1-4}\text{alkyl}$;

R^2 is hydrogen, $\text{C}_{1-4}\text{alkyl}$, or R^1 and R^2 represent $(\text{CH}_2)_{2-4}$ to complete a heterocyclic ring;

R^3 is hydrogen, hydroxyl, $\text{C}_{1-4}\text{alkoxy}$, or halogen;

R^4 and R^5 are independently selected from hydrogen, halogen, nitrile, $\text{C}_{1-4}\text{alkoxy}$,

C₁₋₆alkylthiol, C₁₋₄alkyl, C₁₋₄alkyl substituted with halogen or C₁₋₆alkoxy, or R⁴ and R⁵ represent (CH₂)_m to complete a cycloalkyl ring, or R⁴ and R⁵ together complete a phenyl or thiophene ring which is unsubstituted or substituted with halogen, C₁₋₄alkyl, or C₁₋₄alkoxy;

m = 3 - 4;

5 n = 0 - 2;

R⁶ is hydrogen, hydroxyl, C₁₋₄alkoxy, C₁₋₄alkoxy substituted with hydroxyl, halogen, or NR⁷R⁸, OC(=O)C₁₋₆alkyl, =O, NR⁷R⁸, C₁₋₄alkyl, or C₁₋₄alkyl substituted with hydroxyl, halogen, or NR⁷R⁸, wherein when n = 0, R⁶ is not hydrogen;

X is C₁₋₄alkoxy, hydroxyl or halogen;

10 R⁷ and R⁸ are independently selected from hydrogen, C₁₋₄alkyl, or C(=O)C₁₋₆alkyl;

or pharmaceutically acceptable salts or solvates thereof.

10. The method of claim 8, wherein R¹ and R³ are hydrogen;

R² is C₁₋₄alkyl;

R⁴ and R⁵ are independently selected from halogen, nitrile, C₁₋₄alkoxy, C₁₋₆alkylthiol, C₁₋₄alkyl, C₁₋₄alkyl substituted with halogen, or R⁴ and R⁵ together represent (CH₂)_m to complete a cycloalkyl ring, or R⁴ and R⁵ together complete a phenyl or thiophene ring which is unsubstituted or substituted with halogen, C₁₋₄alkyl;

m = 3 - 4;

n = 1;

20 R⁶ is hydroxyl, C₁₋₄alkoxy, C₁₋₄alkoxy substituted with hydroxyl, halogen, or NR⁷R⁸, OC(=O)C₁₋₆alkyl, NR⁷R⁸, or C₁₋₄alkyl substituted with hydroxyl, halogen, or NR⁷R⁸, wherein when n = 0, R⁶ is not hydrogen;

X is C₁₋₄alkoxy or hydroxyl;

R⁷ and R⁸ are independently selected from hydrogen, C₁₋₄alkyl, or C(=O)C₁₋₆alkyl.

11. The method of claim 8, wherein said compound is:

5-(2-Aminopropyl)-8-bromo-6-methoxy-chroman-3-ol;

5-((R)-2-Aminopropyl)-8-bromo-6-methoxy-chroman-3-ol;

5-(2-Aminopropyl)-6-methoxy-8-methyl-chroman-3-ol;

5 5-((R)-2-Amino-1-hydroxy-propyl)-8-bromo-6-methoxy-chroman-3-ol;

Cyclopropanecarboxylic acid 5-((R)-2-aminopropyl)-8-bromo-6-methoxy-chroman-3-yl ester;

[5-(2-Aminopropyl)-6-methoxy-8-methyl-chroman-3-yl]-methanol;

5-(2-Aminopropyl)-8-iodo-chroman-3,6-diol; or

10 [4-(2-Aminopropyl)-5-methoxy-7-methyl-2,3-dihydro-benzofuran-2-yl]-methanol;

or combinations thereof.

12. A method for the treatment of glaucoma comprising administering a pharmaceutically effective amount of a composition comprising at least one compound of claim 1.

15 13. The method of claim 12, wherein R¹ is hydrogen or C₁₋₄alkyl;

R² is hydrogen, C₁₋₄alkyl, or R¹ and R² represent (CH₂)₂₋₄ to complete a heterocyclic ring;

R³ is hydrogen, hydroxyl, C₁₋₄alkoxy, or halogen;

R⁴ and R⁵ are independently selected from hydrogen, halogen, nitrile, C₁₋₄alkoxy,

C₁₋₆alkylthiol, C₁₋₄alkyl, C₁₋₄alkyl substituted with halogen or C₁₋₆alkoxy, or R⁴ and R⁵

20 represent (CH₂)_m to complete a cycloalkyl ring, or R⁴ and R⁵ together complete a phenyl or thiophene ring which is unsubstituted or substituted with halogen, C₁₋₄alkyl, or C₁₋₄alkoxy;

m = 3 - 4;

n = 0 - 2;

R⁶ is hydrogen, hydroxyl, C₁₋₄alkoxy, C₁₋₄alkoxy substituted with hydroxyl, halogen, or

NR^7R^8 , $\text{OC}(=\text{O})\text{C}_{1-6}\text{alkyl}$, $=\text{O}$, NR^7R^8 , $\text{C}_{1-4}\text{alkyl}$, or $\text{C}_{1-4}\text{alkyl}$ substituted with hydroxyl, halogen, or NR^7R^8 , wherein when $n = 0$, R^6 is not hydrogen;

X is $\text{C}_{1-4}\text{alkoxy}$, hydroxyl or halogen;

R^7 and R^8 are independently selected from hydrogen, $\text{C}_{1-4}\text{alkyl}$, or $\text{C}(=\text{O})\text{C}_{1-6}\text{alkyl}$;

5 or pharmaceutically acceptable salts or solvates thereof.

14. The method of claim 12, wherein R^1 and R^3 are hydrogen;

R^2 is $\text{C}_{1-4}\text{alkyl}$;

R^4 and R^5 are independently selected from halogen, nitrile, $\text{C}_{1-4}\text{alkoxy}$, $\text{C}_{1-6}\text{alkylthiol}$, $\text{C}_{1-4}\text{alkyl}$, $\text{C}_{1-4}\text{alkyl}$ substituted with halogen, or R^4 and R^5 together represent
10 $(\text{CH}_2)_m$ to complete a cycloalkyl ring, or R^4 and R^5 together complete a phenyl or thiophene ring which is unsubstituted or substituted with halogen, $\text{C}_{1-4}\text{alkyl}$;

$m = 3 - 4$;

$n = 1$;

R^6 is hydroxyl, $\text{C}_{1-4}\text{alkoxy}$, $\text{C}_{1-4}\text{alkoxy}$ substituted with hydroxyl, halogen, or
15 NR^7R^8 , $\text{OC}(=\text{O})\text{C}_{1-6}\text{alkyl}$, NR^7R^8 , or $\text{C}_{1-4}\text{alkyl}$ substituted with hydroxyl, halogen, or NR^7R^8 , wherein when $n = 0$, R^6 is not hydrogen;

X is $\text{C}_{1-4}\text{alkoxy}$ or hydroxyl;

R^7 and R^8 are independently selected from hydrogen, $\text{C}_{1-4}\text{alkyl}$, or $\text{C}(=\text{O})\text{C}_{1-6}\text{alkyl}$.

15. The method of claim 12, wherein said compound is:

20 5-(2-Aminopropyl)-8-bromo-6-methoxy-chroman-3-ol;

5-((R)-2-Aminopropyl)-8-bromo-6-methoxy-chroman-3-ol;

5-(2-Aminopropyl)-6-methoxy-8-methyl-chroman-3-ol;

5-((R)-2-Amino-1-hydroxy-propyl)-8-bromo-6-methoxy-chroman-3-ol;

Cyclopropanecarboxylic acid 5-((R)-2-aminopropyl)-8-bromo-6-methoxy-chroman-

3-yl ester;

[5-(2-Aminopropyl)-6-methoxy-8-methyl-chroman-3-yl]-methanol;

5-(2-Aminopropyl)-8-iodo-chroman-3,6-diol; or

[4-(2-Aminopropyl)-5-methoxy-7-methyl-2,3-dihydro-benzofuran-2-yl]-methanol;

5 or combinations thereof.

16. A pharmaceutical composition comprising the compound of claim 1 and at least one carrier.

17. A method to block or bind to serotonin receptors comprising administering an effective amount of at least one compound of claim 1 to a patient.